

Risks and Complications of Posterior Cruciate Ligament Surgery

Ph. Landreau MD
Aspetar: Qatar Orthopaedic and Sports Medicine Hospital


Complications

- In addition to standard risks associated with all orthopedic surgical procedures, posterior cruciate ligament (PCL) reconstruction poses some relatively unique potential complications. These complications arise from a combination of several factors:
 - the relative infrequency of PCL injuries,
 - the lack of knowledge and experience in treating them,
 - the proximity of neurovascular structures to the PCL,
 - and the technically demanding nature of reconstructive procedures

Surgical complications

- Neurovascular injuries
- Osteonecrosis
- Fractures
- Compartmental syndrom


Neurovascular injuries



Zawodny et al.

Neurovascular Injury


- Most feared complication
- Especially in revision surgery
- Popliteal Artery: laceration, occlusion, spasm
- Mostly caused by guide pin and drill during the tibial tibial procedure, sometimes while doing posteromedial approach or capsule manipulation



Zawodny et al.

Neurovascular Injury


- Proximity between artery and nerve and PCL
- Distance 7 - 9 mm between PCL tibial insertion and popliteal artery
- the distance increases significantly to 9 -10 mm at 100° of flexion



Malava MJ, Sethi NS, Totty WG. Proximity of the posterior cruciate ligament insertion to the popliteal artery as a function of the knee flexion angle - implications for posterior cruciate ligament reconstruction. *Arthroscopy* 2000 ; 16 : 796-804

Neurovascular Injury

- Posterior capsule release



Ahn JH, Wang JH, Lee SH, et al. Increasing the distance between the posterior cruciate ligament and the popliteal neurovascular bundle by a limited posterior capsular release during arthroscopic transibial posterior cruciate ligament reconstruction : a cadaveric angiographic study. *Am J Sports Med* 2007 ; 35 : 787-92

How to minimize the NV risk?



Leg position

- 100°
- Tourniquet
- Arthro-pump



Postero-medial approach

- Spinal needle, transillumination
- Thin canula (6 mm)
- Cautious use motorized instruments, arthrosurface and curettes
- Instruments must always be oriented anteriorly
- Limited distal posterior capsule release


Instrumentation

- Specific curved instruments in order to protect the neurovascular area



Tibial tunnel procedure

- Specific guide
- Protection of the vessels
- Cautious drilling under visual control
- 100° of flexion



Tibial tunnel procedure



Fluoroscopy?

Killer Turn

Inlay procedure?

Oliviero et al.

Revision

McAllister et al.

Neurovascular Injury

- Fanelli's trick
 - Posteromedial safety incision

Fanelli GC, Boyd J. How I manage PCL injuries. Oper Tech Sports Med 2009; 17 : 175-83

Osteonecrosis: medial condyle

- Rare
- Less power blood supply than the lateral condyle
- Avoid trauma to the subchondral blood supply by accurate femoral tunnel placement 8 to 10 mm from the articular surface and limit soft tissue dissection
- Double bundle repair: Higher risk?

Athanasian EA, Wickiewicz TL, Warren RF. Osteonecrosis of the femoral condyle after arthroscopic reconstruction of a cruciate ligament: report of two cases. *J Bone Joint Surg Am.* 1995;77:1418-1422.

Reddy AS, Frederick RW. Evaluation of the intraosseous and extraosseous blood supply to the distal femoral condyles. *Am J Sports Med.* 1998;26:415-419.

Fractures

- Tibia or femur
 - tunnels
 - Divergence
 - Hammering staples
- Patella
 - 0.2% to 2.3% incidence for ACL with BTB
 - Bone harvest
 - Quadriceps graft

Zawodny et al.

Voos JE, Drakes MC, Lovich DG, et al. Proximal tibial fracture after anterior cruciate ligament reconstruction using bone-patellar tendon-bone autograft: a case report. *IBS/J.* 2006;4:20-24.

Malik MM, Kunkle KL, Knable KR. Intraoperative complications of arthroscopically assisted ACL reconstruction using patellar tendon autograft. *Arth Course Lect.* 1996;45:297-302.

Compartment Syndrome

- Fluid extravasation into the leg compartments through a rent in the capsule
 - First femoral tunnel, then tibial tunnel

Krysa J, Lofthouse R, Kavanagh G. Gluteal compartment syndrome following posterior cruciate ligament repair. *Injury*. *Int J Care Injured*. 2002;33:835-838.

Conclusion

- Yes, PCL reconstruction is a more dangerous procedure than ACL reconstruction
 - anatomic
 - less experience

