

Hunter 1743

"From Hippocrates to present age, it is universally allowed that ulcerated cartilage is a trouble something, and that, once destroyed, is not repaired"



Cartilage + Cartilage is marvellous: tough, elastic, durable. + If normal, it will last lifetime.

... if damaged, even normal activity will lead to erosion of joint surface. +

Causes of Cartilage injuries

- Sports
- involving torsional forces, high speed sports.
- Trauma - RTA, Bike injuries
- idiopathic - OCD
- Associated with other IA lesions
- Neglected
- Missed
- Ignored




ICRS Arthroscopic **Classification Lesion Thickness** + Grade 0: Normal 1

- + Grade I: Superficial fissures
- + Grade II:<1/2 depth



ICRS Arthroscopic Classification

- Lesion Thickness
- + Grade III: >50% depth but not through subchondral plate.
- Grade IV: lesion through subchondral plate
 - OCD lesions
 - AVN lesions





Treatment options



+ Some light at the end of 'centuries old dark tunnel'?


- Benign neglect
- Debridement Pridie's perforations
- Abrasion Arthroplasty Morselized osteochondral mixture
- Steadmann's Microfracture Periosteal Grafting
- Periosteal Grafting Perichondrial Grafting Osteochondral Allograft Osteochondral Autograft Mosaicplasty ACI and its derivatives Biomaterials


- Biomaterials


Treatment goals





- + Permanent restoration of cartilage surface.
- + Hyaline or Hyaline like regeneration. – Ideal.
- + If not possible, fibrocartilage regeneration.
 - Poor biomechanical properties.
 - Tends to wear off relatively fast.
 - Doest not stand high demand activities.















Hangody's Mosaicplasty

- Cylindrical Osteochondral grafts

 From NWB articular
 - surface.
 - To WB articular surface.
 Indications:
 - Non degenerative
 - 1.5 to 4 cm²
 - Isolated
- Absolute contraindications:
- Age > 50
 Size > 4-8 cm²
- Size > 4-8 cm²
 Depth > 10 cm.












- Kevin Stone's morsellised autogenous osteochondral mixture.
- + Periosteal grafting.
- + Osteochondral autogenous transfer.



Treatment Protocol....Algorithm

- + Grade 1: Conservative
- Grade 2:
- Debridement
- Extensive controlled rehabilitation
- + Grade 3 & 4:
 - < 1cm: Benign neglect or microfracures</p>
 - 1-2 cm: Microfractures
 - 1.5 to 4 cm: Mosaicplasty
 - 3 to 10 cm: ACI, osteochondral allografts




Manual medial medial medial medial medial menti menti menti menti menti menti menti

Measurement of the defect size: 20 11mm















... still a mountain to climb...









